

Acetaminophen/Ibuprofen Dosage Chart

Acetaminophen (Tylenol® or another brand)

Give every 6 hours as needed. (Also available in suppositories; use the same number of mg.)

*Do not give more than 4 doses in 24 hours.

Weight in pounds (lbs.)	Elixir 1 teaspoon = 160 mg/5ml	Chewable 1 tablet = 80 mg	Jr. strength 1 caplet = 160 mg	Reg. Strength 1 tablet = 325mg
6-11 lbs.	1/4 teaspoon(1.25 ml)			
12-17 lbs.	½ teaspoon(2.5 ml)			
18-23 lbs.	³ / ₄ teaspoon(3.75 ml)			
24-35 lbs.	1 teaspoon(5 ml)	2 tablets		
36-47 lbs.	1½ teaspoons(7.5 ml)	3 tablets		
48-59 lbs.	2 teaspoons(10 ml)	4 tablets	2 caplets	1 tablet
60-71 lbs.	2½ teaspoons(12.5 ml)	5 tablets	2½ caplets	1 tablet
72-95 lbs.	3 teaspoons(15 ml)	6 tablets	3 caplets	1½ tablets
96+ lbs.			4 caplets	2 tablets

Ibuprofen (Advil, Motrin, or another brand)

Give every 6 to 8 hours as needed; **always** with food. Other strengths available by prescription.

*Do not give more than 4 doses in 24 hours.

Weight in pounds (lbs)	Infant drops (40mg/1 ml)	Children's liquid 1 teaspoon = 100 mg/5ml	Chewable Tablets 1 tablet = 100 mg	Tablets 1 tablet = 200 mg
11-21 lbs.	¹ / ₄ teaspoon (1.25ml)	½ teaspoon(2.5 ml)	½ tablet	
22-32 lbs.	½ teaspoon (2.5ml)	1 teaspoon(5 ml)	1 tablet	
33-43 lbs.		1½ teaspoons(7.5 ml)	1 ½ tablets	
44-54 lbs.		2 teaspoons(10 ml)	2 tablets	1 tablet
55-65 lbs.		2½ teaspoons(12.5 ml)	2 ½ tablets	
66-87 lbs.		3 teaspoons(15 ml)	3 tablets	1½ tablets
88+ lbs.		4 teaspoons(20 ml)	4 tablets	2 tablets

Questions?

This sheet is not specific to your child but provides general information. If you have any questions, please call your clinic or pharmacy.